

# Understanding possible side effects with VYLOY<sup>®</sup>

Use this guide to learn more about  
common side effects that may occur  
during or after your VYLOY infusion

## What is VYLOY?

VYLOY (zolbetuximab-clzb) is a prescription medicine used to treat adults with cancer of the stomach (gastric cancer) or cancer located where the esophagus joins the stomach (gastroesophageal junction cancer). VYLOY is used in combination with chemotherapy that contains fluoropyrimidine and platinum as the first treatment when your gastric or gastroesophageal junction cancer:

- cannot be removed with surgery or has spread to other parts of the body,
- is HER2-negative, **and**
- your tumor tests positive for “claudin (CLDN)18.2.”

It is not known if VYLOY is safe and effective in children.

## IMPORTANT SAFETY INFORMATION

**Before receiving VYLOY, tell your healthcare provider about all of your medical conditions, including if you:**

- have nausea or vomiting.
- are pregnant or plan to become pregnant. It is not known if VYLOY will harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if VYLOY passes into your breast milk. Do not breastfeed during treatment with VYLOY and for 8 months after the last dose.

**Tell your healthcare provider about all the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Please see additional [Important Safety Information](#) on page 4, and read the [Patient Information](#).

## WHAT POTENTIAL SIDE EFFECTS CAN OCCUR DURING TREATMENT WITH VYLOY?

**VYLOY (zolbetuximab-clzb) may cause side effects in some people, but knowing what to expect can help you be prepared**

Side effects can occur with treatment. It is important to keep track of how you feel during and after your VYLOY infusion. Be sure to share any questions and concerns with your healthcare team.

**VYLOY may cause serious side effects, including:**



### Allergic reactions, including anaphylaxis and infusion-related reactions

- Allergic reactions are common during treatment with VYLOY and can sometimes be serious
- Serious allergic reactions can happen during or after your VYLOY infusion, including life-threatening allergic reactions, and serious infusion-related reactions that may lead to death
- Your healthcare provider will monitor you during your infusion and for 2 hours after or longer if needed
- Tell your healthcare provider or get emergency medical help right away if you get any of the following symptoms of a serious allergic reaction during or after your infusion of VYLOY:
  - itchy, raised bumps on the skin (hives)
  - coughing that does not go away
  - nausea or vomiting
  - stomach (abdominal) pain
  - increased saliva
  - breathing problems such as wheezing
  - throat tightness or change in voice
  - fever
  - chest discomfort
  - chills or shaking
  - back pain



### Severe nausea and vomiting

- Nausea and vomiting are common during treatment with VYLOY and can sometimes be severe
- Nausea and vomiting happened more often during the first treatment cycle
- Before you receive each VYLOY infusion, your healthcare provider will give you medicines to help prevent nausea and vomiting
- Tell your healthcare provider right away if nausea or vomiting does not go away or gets worse



### Keep your healthcare team informed

Speak up if you experience any new side effects, ones that do not go away, or if they get worse.

If you have trouble tolerating treatment, your treatment may be adjusted accordingly.

# WHAT POTENTIAL SIDE EFFECTS CAN OCCUR DURING TREATMENT WITH VYLOY? (CONT.)

3

The most common side effects of VYLOY include:



## Stomach problems

- decreased appetite
- diarrhea
- stomach (abdominal) pain
- constipation
- decreased weight




## Changes in lab tests

- decreased white blood cells, red blood cells and platelets
- decreased protein (albumin) in the blood
- changes in kidney function tests
- changes in blood sugar (glucose)
- changes in liver function tests
- changes in body salts (electrolytes) in your blood



## Other reactions

- tiredness
- fever
- tingling or numbness of the arms and legs



Your healthcare provider may slow the rate of your infusion, temporarily stop, or completely stop treatment with VYLOY if you have certain side effects.

These are not all of the possible side effects of VYLOY.

**Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch)**



Here are some tips that have helped others:

- Keep a **treatment journal** to share with your team
- Stay **hydrated**. Drink plenty of water and other fluids
- Work with your healthcare team to make a plan for **good nutrition**
- Try to **focus on the positive**. Surround yourself with people and things you love
- Take a **walk or sit outside** to get some fresh air

Please see additional [Important Safety Information](#) on page 4, and read the [Patient Information](#).



For more information about how to prepare for your VYLOY infusion, visit [VYLOY.com](https://www.vyloy.com)

## IMPORTANT SAFETY INFORMATION (cont'd)

### What are the possible side effects of VYLOY (zolbetuximab-clzb)?

VYLOY may cause serious side effects, including:

- **Allergic reactions, including anaphylaxis and infusion-related reactions.** Allergic reactions are common during treatment with VYLOY and can sometimes be serious. Serious allergic reactions can happen during or after your VYLOY infusion, including life-threatening allergic reactions and serious infusion-related reactions that may lead to death. Your healthcare provider will monitor you during your infusion and for 2 hours after or longer if needed. Tell your healthcare provider or get emergency medical help right away if you get any of the following symptoms of a serious allergic reaction during or after your infusion of VYLOY:
  - itchy, raised bumps on the skin (hives)
  - coughing that does not go away
  - nausea or vomiting
  - stomach (abdominal) pain
  - increased saliva
  - breathing problems such as wheezing
  - throat tightness or change in voice
  - fever
  - chest discomfort
  - chills or shaking
  - back pain
- **Severe nausea and vomiting.** Nausea and vomiting are common during treatment with VYLOY and can sometimes be severe. Nausea and vomiting happened more often during the first treatment cycle. Before you receive each VYLOY infusion, your healthcare provider will give you medicines to help prevent nausea and vomiting. Tell your healthcare provider right away if nausea or vomiting does not go away or gets worse.

### The most common side effects of VYLOY include:

- tiredness
- decreased appetite
- diarrhea
- tingling or numbness of the arms or legs
- stomach (abdominal) pain
- constipation
- decreased weight
- fever
- decreased white blood cells, red blood cells and platelets
- decreased protein (albumin) in the blood
- changes in kidney function tests
- changes in blood sugar (glucose)
- changes in liver function tests
- changes in body salts (electrolytes) in your blood

Your healthcare provider may slow the rate of your infusion, temporarily stop, or completely stop treatment with VYLOY if you have certain side effects. These are not all of the possible side effects of VYLOY. Call your doctor for medical advice about side effects. You may report side effects to the FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](https://www.fda.gov/medwatch).



Please read the [Patient Information](#).



© 2025 Astellas Pharma Inc. or its affiliates.  
All rights reserved. MAT-US-VYL-2025-00187 04/25  
All trademarks are the property of their respective owners.